

## Role of Autologous Platelet Rich Plasma in Scar Management

Adithya Kevin J<sup>1</sup>, Ravi Kumar Chittoria<sup>2\*</sup>, Nishad Kerakada<sup>3</sup>, Neljo Thomas<sup>3</sup>, Barathkumar Singh.P<sup>3</sup>,  
Jacob Antony Chakiath<sup>3</sup>, P. Chakkravarthy<sup>1</sup>, Varun<sup>1</sup>

<sup>1</sup>Department of General Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Pondicherry, India

<sup>2</sup>Department of Plastic Surgery & Telemedicine, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Pondicherry, India

<sup>3</sup>Department of Plastic Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Pondicherry, India

### Research

Please cite this paper as: Kevin JA, Chittoria RK, Kerakada N, Thomas N, Singh BP, Chakiath JA, Chakkravarthy P, Varun. Role of Autologous Platelet Rich Plasma in Scar Management. AMJ 2024;17(3):1198-1200.  
<https://doi.org/10.21767/AMJ.2024.4026>

#### Corresponding Author:

Ravi Kumar Chittoria  
Department of Plastic Surgery & Telemedicine,  
Jawaharlal Institute of Postgraduate Medical Education and Research(JIPMER),  
Pondicherry, India  
Email id: drchittoria@yahoo.com

### Abstract

Autologous platelet rich plasma (APRP) has gained its importance in medical field since its first use in sports medicine and open-heart surgeries. It is widely used in plastic surgery and in cosmetic medicine because of its wound healing properties. Scar management is the most important part of cosmesis and is an important factor in improving the post traumatic quality of life of the patient. Here we present the case of a 17-year-old male with a post traumatic wound on the right knee treated with APRP for scar management

**Key Words:** Autologous platelet rich plasma (APRP), scar management.

### Introduction

Scar management represent an important field in medical science as scars can negatively impact the quality of life of an individual. In addition to the cosmetic blemish, pain, tenderness, itchinnessincludes few other problems associated with a scar. A scar located in a precarious

position can cause functional limitation in the form of scar contraction. Scar aesthetics can also have a negative influence on psychosocial health. This warrants the importance of providing a scar that is aesthetically acceptable and has minimal functional impairment. A number of modalities have been tried and tested over time to achieve the above goal that include fractional CO2 laser therapy<sup>1,2</sup>. Autologous platelet rich plasma (APRP) is a safe, easily accessible and upcoming modality that is finding its use in various fields of medicine. Its use in scar management is still being studied but is an application worth concentrating.

### Materials and Methods

This study was carried out in the department of Plastic Surgery in a tertiary care centre in South India after getting written informed consent from the patient and approval from the department. A 17-year-old male with no known comorbidities presented with an alleged history of road traffic accident (RTA) causing him to sustain fracture to the right tibia with associated avulsion injury resulting in a raw area over the right knee. The tibial fracture had been treated with closed reduction with external fixation leaving the raw area on the right knee. that was treated with Hydrojet wound debridement withprolotherapy, insulin therapy and negative pressure wound therapy for wound bed preparation followed by keystone flap for closure. This resulted in a scar formation. The initial Vancouver scar scale was calculated to be 9 (Figure 1). Autologous platelet rich plasma (APRP) obtained by standard double centrifugation protocol using 10cc of the patient's blood was used for reducing postoperative scar formation with APRP being injected over 8 sittings on weekly basis spanning over 2 months (Figure 2). The scar being assessed with Vancouver scar scale and patient satisfaction.

## Results

The scar score improved with adequate patient satisfaction with a final Vancouver scar scale of 6 (Figure 3). No complications were noted. The therapy was well tolerated

## Discussion

Autologous platelet rich plasma (APRP) as the name implies refers to the plasma derived from the patient's own blood with a platelet count higher than the platelet count in the peripheral blood of the patient. Historically having been used to treat thrombocytopenia, the use in other specialities became widespread with its use in sports medicine to treat musculoskeletal injuries. Its use in wound management results from the observation that wounds have a proinflammatory environment that impairs healing. In addition, wounds have a high protease activity that impairs functioning of growth factors. APRP used in a chronic wound serves as a source of growth factors and thence has mitogenic, angiogenic and chemotactic properties. APRP has also been shown to stimulate human dermal fibroblast proliferation and thus increasing the deposition of TYPE I collagen, the above mechanism being proposed to its use in scar management<sup>3-5</sup>. Application of activated APRP also provides 5 to 10 times the normal concentration of growth factors that include PDGF, VEGF, TGF- $\beta$  locally also accelerating wound healing. Addition of calcium salts also helps in activation of platelets<sup>6-9</sup>.

Usually, around 1 to 1.5 ml of APRP can be obtained from 10 ml of patient's blood. Hence, the disadvantage of the use of APRP lies in its use in wounds of a large surface area that would require a large volume of blood which in a patient with a chronic non healing wound or a traumatic wound requires consideration. Moreover, injecting APRP prior to grafting or flap placement could provide an uneven surface for a regular uptake.

## Conclusion

Autologous platelet rich plasma is an effective measure in improving scar remodelling and is a good choice for treating scars provided the patient has a good functional status and surface area to be treated is small.

## References

1. Rubina Alves, Ramon Grimalt. A review of platelet-rich plasma: History, biology, mechanism of action, and classification. *Skin appendages discord*. 2018;4:18-24. Doi:<https://karger.com/sad/collection/25/Dermatolog>  
y

2. Makki M, Younes AE, Fathy A, et al. Efficacy of platelet-rich plasma plus fractional carbon dioxide laser in treating posttraumatic scars. *Dermatologic Therapy*. 2019;32(5):e13031. Doi: <https://doi.org/10.1111/dth.13031>
3. Mohan PB, Gupta S, Chittoria RK, et al. Autologous platelet-rich plasma enriched pixel grafting. *J Cutan Aesthet Surg*. 2020;13(4):340-3. Doi: 10.4103/JCAS.JCAS\_112\_19
4. Hersant B, SidAhmed-Mezi M, Bosc R, et al. Autologous platelet-rich plasma/thrombin gel combined with split-thickness skin graft to manage postinfectious skin defects: A randomized controlled study. *Adv Skin Wound Care*. 2017;30(11):502-8.
5. Cohn CS, Lockhart E. Autologous platelet-rich plasma: evidence for clinical use. *Current opinion in hematology*. 2015;22(6):527-32. Doi: 10.1097/MOH.0000000000000183
6. Sudhanva HK, Panday S, chittoria RK, et al. Role of APRP in the successful uptake of Split Skin Graft. *Dermatology international*. 2016(1)2.
7. Elankumar S, Sudhanva HK, Abhinav A, Chittoria RK. APRP spray devices: A novel technique of applying APRP. *Dermatology international*. 2017;2(2).
8. Weibrich G, Kleis WK, Hafner G, et al. Growth factor levels in platelet-rich plasma and correlations with donor age, sex, and platelet count. *J Craniomaxillofac Surg*. 2002;30(2):97-102. Doi: <https://doi.org/10.1054/jcms.2002.0285>
9. Yuan T, Zhang CQ, Tang MJ, et al. Autologous platelet-rich plasma enhances healing of chronic wounds. *Wounds*. 2009;21(10):280-5.

## CONFLICTS OF INTEREST

None

## DECLARATIONS

### Authors' contributions

All authors made contributions to the article

### Availability of data and materials

Not applicable

### Financial support and sponsorship

None

## CONSENT FOR PUBLICATION

Not applicable

## Figures



**Figure 1: Initial status of the scar on the anteromedial aspect of the right knee (Vancouver scar scale 9)**



**Figure 2: APRP injection into the scar**



**Figure 3: Final result of the scar after completion of treatment with APRP (Vancouver scar scale 6).**